

From the two-dimensional T_H1 and T_H2 phenotypes to high-dimensional models for gene regulation

Henk-Jan van den Ham and Rob J. de Boer

Theoretical Biology and Bioinformatics, Universiteit Utrecht, Utrecht, The Netherlands

Keywords: helper T cell differentiation, master regulator, mathematical model

Abstract

The T_H1/T_H2 paradigm has been used for decades to characterize phenotypically different immune responses. Recent discoveries, e.g. T_H17 cells are adding more dimensions to the helper T cell framework, and the T_H1/T_H2 paradigm is currently being extended to include these new phenotypes. Previous mathematical models cannot easily be extended to accommodate these new phenotypes, and therefore these discoveries call for a new type of models. We devised a new model of helper T cell differentiation that describes expression of, and interactions between, the master regulators determining the phenotypic polarization of helper T cells. The model is able to describe any number of master regulators and is grounded on transcription factors binding promoter sites and binding each other. The model allows for stable switches between several different phenotypes. Furthermore, the model accounts for the kinetics of FoxP3 and GATA3 mRNA expression measured after stimulating naive helper (CD4+CD45RA+) T cells under various circumstances. Due to its n -dimensional character, this model may easily be applied to other developmental processes that involve master regulators.

Introduction

Helper T cells are CD4+ T lymphocytes that have an important role in determining the nature of the adaptive immune response. Since the 1980s, helper T cells have been classified in two major types, namely the T_H1 or T_H2 phenotype (1). T_H1 cells promote cellular immunity, which is associated with anti-viral responses and tumour surveillance, whereas T_H2 cells promote humoral responses to extracellular parasites and are involved in allergies. At a molecular level, the differentiation of a naive helper T cell towards either the T_H1 or T_H2 phenotype is associated with the expression of phenotype-specific transcription factors Tbet (2) and GATA3 (3), respectively. These two mutually exclusive transcription factors are 'both necessary and sufficient' for inducing a particular cellular phenotype and have therefore given the name 'master regulators' (4) to underscore their importance in cell differentiation.

The two-dimensional T_H1/T_H2 paradigm has been a very successful foundation of immunology for the past 20 years. However, the recent discovery of T_H17 cells confirmed earlier evidence that helper T cells may adopt phenotypes other than T_H1 and T_H2 (5). T_H17 cells are characterized by the production of IL-17 that drives rapid neutrophil recruitment in response to bacterial and fungal infections (6, 7) and is thought to play a role in various autoimmune disorders (8).

In mice, naive helper T cell differentiation towards the T_H17 phenotype is induced by a combination of the cytokines IL-6 and transforming growth factor (TGF) β (9), mediated by master regulators ROR γ t (10) and ROR α (11). The induction of this phenotype in humans is different, but the resulting T cell phenotype appears to be similar (12, 13). Interestingly, TGF β stimulation alone induces expression of another master regulator, FoxP3, which prompts the cell to adopt a regulatory (Treg) phenotype (14–16). Therefore, the T_H17 phenotype does not only provide convincing evidence for the existence of more than two helper T phenotypes but also links the fairly separate fields of T_H1/T_H2 and Treg immunology to each other. As a result, the T_H1/T_H2 paradigm is currently being extended into a framework that includes at least T_H1 , T_H2 , T_H17 and Treg cells (17), which functionally correspond to promoting the cellular, humoral, rapid anti-bacterial/anti-fungal and anti-inflammatory response, respectively.

To obtain more insight into helper T cell differentiation at a molecular level, mathematical models have been employed to describe the expression of the master regulators that control the different phenotypes (18–21). These models have so far only described the T_H1 and T_H2 phenotypes and do not accommodate newly identified lineages

easily. We aim to extend these studies by formulating a more general framework for describing master regulators. Using the molecular biology of transcription factors, involving promoter binding and complex formation, we develop a mathematical model that intends to describe the cardinal intracellular dynamics of any number of master regulators, so that novel regulators can be incorporated as they are discovered. Furthermore, we show that the model is in agreement with previously observed master regulator kinetics in human naive CD4+CD45RA+ T cells stimulated under T_h2 - and Treg-inducing conditions (22).

Methods

Mathematical model

Master regulators are initially up-regulated by external activators (E), inducing the binding of transcription factors mediating cytokine stimulation to the promoter ($P_0 \rightarrow P_E$). Additionally, master regulator protein dimers (D) bind to the promoter ($P_0 \rightarrow P_D$), boosting the transcription of its own mRNA (R), which in turn is translated into master regulator monomers (M). Protein monomers may be sequestered into complexes with other master regulators (N), thereby preventing their dimerization into homodimers. In principle, sequestered master regulator monomers can dissociate from the complex, but if the dissociation constant is small compared with the degradation constant of the complex, this hardly affects the results (data not shown). Master regulators in the complex are therefore taken to be directly degraded without the possibility of dissociation. This system (depicted schematically in Fig. 1) may be translated into a set of simple ordinary differential equations (Equations 1–5) using mass action kinetics:

$$\frac{dR}{dt} = c(P_D + P_E) - d_r R \quad (1)$$

$$\frac{dM}{dt} = lR - d_p M - 2k_1 M^2 + 2k_{-1} D - bMN \quad (2)$$

$$\frac{dD}{dt} = k_1 M^2 - k_{-1} D - d_p D \quad (3)$$

$$\frac{dP_D}{dt} = k_2(1 - P_D - P_E)D - k_{-2} P_D \quad (4)$$

$$\frac{dP_E}{dt} = k_3(1 - P_D - P_E)E - k_{-3} P_E \quad (5)$$

where c and l are transcription and translation rates, d_r and d_p are mRNA and protein degradation rates, k_1 , k_2 and k_3 are association rates, k_{-1} , k_{-2} and k_{-3} are dissociation rates and b is the rate of complex formation between master regulator protein monomers.

Using quasi-steady-state assumptions and the ‘conservation equation’ $P_0 + P_D + P_E = 1$ (that bounds the promoter ‘activity’ between zero and one), these five equations can be simplified to only one equation for the expression of the master regulator. Assuming that the binding of dimers and external activators to the promoter is a fast process (i.e. $\frac{dP_D}{dt} = \frac{dP_E}{dt} = 0$), we obtain a multiple substrate Michaelis–Menten




Fig. 1. Outline of master regulator gene regulation, which leads to key master regulator properties external activation, self-activation and inhibition. The master regulator gene is initially activated by external activators, for instance, in response to cytokine stimulation. The master regulator protein monomers are produced in response to the external activators and these bind as dimers to the master regulator promoter. Additionally, the master regulator monomers can be sequestered into non-active complexes by binding to monomers from other master regulator genes. For mathematical formulation of the model, please see the Methods.

term as a function for the relative activity of the promoter in terms of master regulator dimers D and external activators E :

$$P = P_D + P_E = \frac{\frac{k_2}{k_{-2}} D + \frac{k_3}{k_{-3}} E}{1 + \frac{k_2}{k_{-2}} D + \frac{k_3}{k_{-3}} E}$$

Proceeding with this expression and making quasi-steady-state assumptions for mRNA and dimers (i.e. $\frac{dR}{dt} = \frac{dD}{dt} = 0$), substitution into Equation (2) yields a single differential equation describing the expression level of the master regulator.

$$\frac{dM}{dt} = \frac{cl}{d_r} \left[\frac{\frac{k_2}{k_{-2}} D + \frac{k_3}{k_{-3}} E}{1 + \frac{k_2}{k_{-2}} D + \frac{k_3}{k_{-3}} E} \right] - d_p M - 2k_1 M^2 + 2k_{-1} \frac{k_1 M^2}{k_{-1} + d_p} - bMN$$

Rescaling with respect to time ($\tau = d_p t$) and introducing new parameters, we obtain

$$\frac{dM}{d\tau} = \rho \left[\frac{\alpha M^2 + \varepsilon}{1 + \alpha M^2 + \varepsilon} \right] - M(1 + \mu M + \beta N) \quad (6)$$

with five parameters describing ‘gene production’ $\rho = \frac{cl}{d_r d_p}$, ‘homodimerization’ $\mu = \frac{2k_1}{k_{-1} + d_p}$, ‘promoter affinity’ $\alpha = \frac{k_2}{2k_{-2}}$, ‘external activation’ $\varepsilon = \frac{k_3}{k_{-3}} E$ and ‘inhibition’ $\beta = \frac{b}{d_p}$, respectively (Box 1).

Data and fitting

The data used to fit the model were obtained from Mantel *et al.* (22) (Figs 4 and 5). Values were extracted using Plot Digitizer, a tool to quantify information from graphical plots. The obtained values were normalized to the maximum expression of FoxP3 and GATA3, respectively, and were fit manually using GRIND, the GReat INtegrator of Differential equations. All data sets were fit simultaneously, i.e. only the

Box 1. Mathematical terminology

Steady states: When a cell is in a steady state, it remains in this state until it is forced out by some external stimulus. The steady state is said to be stable if the cell returns to the steady state after a small transient perturbation; if a cell does not return, this is a non-stable steady state.

Vector field: When a cell is not in a steady state, but changes over time, the vector field shows the direction in which the concentrations of master regulators in the cell will change. The vector field can be used to determine the stability of steady states.

Basin of attraction: Every stable steady state has basin of attraction, or 'catchment area' around it, composed of the states that would approach this steady state.

Separatrix: When there are several stable steady states in the system, each has a basin of attraction. A separatrix is the border between these two basins or the 'watershed' between 'catchment areas'.

Hysteresis: When a cell is transiently activated into a differentiation pathway, it can remain in this differentiated state after the original stimulus has been removed. This memory for a previously adopted state is known as hysteresis [see (23)].

Bifurcation: Here, a bifurcation is a change in a steady state of a system. States can appear and disappear at so-called saddle-node bifurcation points. A bifurcation analysis is summarized in a diagram depicting the steady states that are present for a range of parameter values. This parameter is known as the bifurcation parameter.

external activation parameters ε_1 and ε_2 were allowed to be different between data sets, while all other parameters were kept constant. The gene production parameter ρ was estimated from the data by observing that the average expressions of FoxP3 and GATA3 increased roughly 4-fold between 0 and 96 h. Initial values were set to measurement values at $t = 0$.

Results*Developing a new model*

A naive helper T cell is activated when its TCRs recognize antigen mounted on the MHC of a professional antigen-presenting cell. In response to particular co-stimulation and the local cytokine environment, the cell ultimately adopts one of several distinct helper T phenotypes (7, 24). In order to adopt the T_{h2} phenotype, a helper T cell needs to express high levels of the master regulator GATA3. GATA3 gene is up-regulated by transcription factors induced by TCR signalling and co-stimulation, such as nuclear factor κ B (NF κ B) and nuclear factor of activated T cells (NFAT) (25, 26). GATA3 is also up-regulated by specific cytokine signals, via the IL-4 receptor and Stat6, which is part of the Jak-Stat signalling pathway (27). Finally, GATA3 dimerizes and promotes its

own transcription in a Stat6-independent manner (28). Likewise, the T_{h1} phenotype requires high expression of Tbet, which is self-activating via the extracellular cytokine route, and is also up-regulated by NF κ B and NFAT. Its specific inducing cytokine is IL-12, which signals via Stat1 and Stat4 (3, 29, 30).

Consistent with the opposite phenotypes they induce, GATA3 and Tbet are typically not stably expressed in the same helper T cell. Three mechanisms have been described to accomplish mutual exclusion. Firstly, both GATA3 and Tbet repress oppositely polarizing cytokines, effectively blocking the inducing signals of the opposite phenotype (31). Secondly, up-regulation of either gene leads to epigenetic modification of the other locus, resulting in decreased transcription (31). Lastly, the TEC family kinase inducible T cell kinase (ITK) mediates the binding of GATA3 and Tbet in the nucleus forming a complex (32, 33). The GATA3–Tbet complex prevents both GATA3 and Tbet from binding to their promoter and thereby interferes with the self-activation mechanism of these master regulators. Since the last mentioned is a clear intracellular mechanistic interaction, this binding mechanism will be used in our model to describe the intracellular crosstalk between master regulators.

Using the general properties of GATA3–Tbet regulation as a paradigm, we formulate a mathematical 'master regulator' model (depicted schematically in Fig. 1) including (i) up-regulation upon T cell activation and cytokine stimulation, (ii) self-activation of their own transcription (i.e. positive feedback) and (iii) inhibitory interactions with other master regulators. Using quasi-steady-state assumptions, mathematical formulation of these mechanisms (see Methods) yields the following model:

$$\frac{dM}{d\tau} = \rho \left[\frac{\alpha M^2 + \varepsilon}{1 + \alpha M^2 + \varepsilon} \right] - M(1 + \mu M + \beta N) \quad (7)$$

where M is the scaled concentration of the master regulator in the cell. A single production term, $\rho \left[\frac{\alpha M^2 + \varepsilon}{1 + \alpha M^2 + \varepsilon} \right]$, and a single degradation term, $M(1 + \mu M + \beta N)$, can be identified (Fig. 2). The production term (derived in the Methods) is a sigmoid Michaelis–Menten equation. The equation has five scaled parameters, consisting of gene production ρ , homodimerization μ , promoter affinity α , external activation ε and inhibition β .

Although based on mechanistic interactions, this remains a caricature model that simplifies reality: many of the details concerning activation of and interactions between master regulators are absent from the model. Potentially important post-translational modifications, such as phosphorylation, are not included because this would make the model much more complex. Such a complex model would have too many unknown parameters, and hence be less realistic, and would be difficult to understand (34). This will nevertheless be a good model if it provides new insights and if it proves to be extendable with future data.

A simple master regulator

The master regulator model describes master regulator dynamics by explicitly stating the contribution of cellular processes towards a change in concentration. Disregarding the interactions with other master regulator genes, the master




Fig. 2. Steady states of a system with one master regulator. The production term, $\rho \left[\frac{\alpha M^2 + \epsilon}{1 + 2\alpha M^2 + \epsilon} \right]$ (solid line), and the degradation term, $M(1 + \mu M)$ (dashed line), are represented as functions of the master regulator monomer concentration M . Solid and open squares represent stable and non-stable steady states, respectively. Approximations for the steady state expression level of the master regulator in terms of the parameters are $M \approx \frac{\rho\epsilon}{1+\epsilon}$, $M \approx \frac{1}{\rho\alpha - \mu} - \rho\epsilon$ and $M \approx \frac{1}{2\alpha}(\sqrt{1 + 4\rho\mu} - 1)$ for states 0, 1 and 2, respectively (see Methods for details).

regulator concentration only depends on the production and degradation of gene products. Due to the feedback of the master regulator on itself, the system is expected to go to equilibrium, i.e. a master regulator concentration where production and degradation of gene products are equal. Plotting production and degradation as a function of the master regulator concentration shows that there are three concentrations where production equals degradation (Fig. 2). These ‘steady states’ have been numbered 0, 1 and 2 in accordance with the expression level of the master regulator. State 0 is a resting, or naive, state with low expression of the master regulator. Once the master regulator is activated to high expression, the system approaches state 2, which is therefore the ‘active’ state. State 1 is not stable and separates the two stable states: it acts as a ‘separatrix’ enabling switches between the two stable states. The sigmoid shape of production term (Fig. 2) is the direct result of dimerization; dimerization is therefore critical to obtain naive and active states within this master regulator model.

The master regulator model describes master regulator dynamics in terms of a number of cellular processes, namely gene production ρ , self-activation α , homodimerization μ and external activation ϵ . Using the parameters that represent these cellular processes as ‘bifurcation parameters’, we observe that all parameters have a bistable range, i.e. a region where both the resting and active state of the master regulator are present (Fig. 3). This means that changes in any of these cellular processes can eliminate steady states by a ‘(saddle-node) bifurcation’, and let the system approach other stable steady states. Changes in gene production (ρ), self-activation (α) and dimerization (μ) (Fig. 3a, c, and d) can all lead to two different bifurcations, and let a cell switch from its resting state 0 to its active state 2 and vice versa. However, changing the external activation (ϵ) (Fig. 3b) only leads to one possible bifurcation that eliminates the naive




Fig. 3. Bifurcation analysis of the single master regulator model for the parameters gene production ρ , external activation ϵ , homodimerization μ and promoter affinity α . The bold and thin lines represent stable and non-stable steady states, respectively (see Box 1). The identity of the state is indicated by state numbers. The boxes represent (saddle-node) bifurcations at which states can appear or disappear, and arrows indicate the direction of the vector field. Default parameter setting: $\rho = 5$, $\epsilon = 0.01$, $\mu = \alpha = 1$ and $\beta = 0$.

state 0. Increasing external activation can therefore lead to a shift from the naive state to the active state 2, but after this state has been approached, decreasing external activation will not cause the cell to go back to the naive state. By triggering the self-activation process, a transient increase in external activation may act as a permanent switch that changes the master regulator expression level from low to high.

The concentrations of the master regulator in the different steady states obviously depend on the cellular processes. Using a conventional Taylor expansion (see Methods), approximations of the steady states were calculated (see legend of Fig. 2). The level of master regulator expression in the resting state 0 is determined primarily by the external activation rate ϵ . The level of expression in the active state 2 primarily depends on the gene production rate ρ and homodimerization rate μ , since state 2 is located near the maximum of the production curve (Fig. 2).

Two master regulators: T_{h1} and T_{h2}

Helper T phenotypes have been shown to depend crucially on the interaction between multiple master regulators. Using the original T_{h1}/T_{h2} paradigm as an illustration, we investigate a system of two master regulators. Introducing a subscript to denote different master regulators and their parameters, one can formulate a two-dimensional system as

$$\frac{dM_1}{d\tau} = \rho \left[\frac{\alpha_1 M_1^2 + \varepsilon_1}{1 + \alpha_1 M_1^2 + \varepsilon_1} \right] - M_1(1 + \mu_1 M_1 + \beta M_2) \quad (8)$$

$$\frac{dM_2}{d\tau} = \rho \left[\frac{\alpha_2 M_2^2 + \varepsilon_2}{1 + \alpha_2 M_2^2 + \varepsilon_2} \right] - M_2(1 + \mu_2 M_2 + \beta M_1) \quad (9)$$

The gene production parameter ρ determines the maximal production rate of a master regulator in a cell and depends on the transcription and translation rates of the cellular machinery. In order to have both naive and polarized states, the production rate ρ should be sufficiently high (Fig. 3a) and is taken to be the same for both genes within the cell.

Phase plane analysis of Equations (8) and (9) (with in total eight parameters) shows that this system has nine different steady states, of which four are stable (Fig. 4a and b). The stable points represent the possible differentiation states that

a helper T cell can attain. When Tbet and GATA3 are both lowly expressed, the T cell is naive (state 00) and resting. When either Tbet or GATA3 is highly expressed, the T cell adopts a T_{h1} (state 20) or a T_{h2} phenotype (state 02), respectively. When both Tbet and GATA3 are highly expressed (state 22), the T cell is active but unpolarized. Due to the mutual inhibition between Tbet and GATA3, this high co-expression normally does not occur in helper T cells, and we have searched for parameter values where the stable high co-expression state (22) is absent from the model. Bifurcation analysis of the inhibition parameter β shows that one can indeed choose an inhibition rate β such that state 22 is non-stable (Fig. 4a and c) or absent (Fig. 4a and d), leaving the naive, T_{h1} and T_{h2} phenotypes as the three stable states of the model.

Helper T cell expression of Tbet and GATA3 is induced by external signals, such as T cell activation and cytokine signalling, thereby skewing the cell towards a certain




Fig. 4. Bifurcation analysis of the inhibition parameter β in a T_{h1}/T_{h2} (two-dimensional) system. The analysis shows a (pitchfork) bifurcation destabilizing the high co-expression state (22). Three phase planes show the nullclines for the indicated parameter values. The states of the system are indicated by state numbers, and stable and non-stable states are indicated by heavy and light lines (bifurcation diagrams) or by solid and open squares (phase plane diagrams). This is a symmetric system with parameter values: $\rho = 5$, $\mu_1 = \mu_2 = \alpha_1 = \alpha_2 = 1$, $\varepsilon_1 = \varepsilon_2 = 0.01$. We display phase plane diagrams for (b) $\beta = 0.5$, (c) $\beta = 1$ or (d) $\beta = 1.5$.

phenotype. This is modelled by parameters ε_1 and ε_2 for T_{h1} and T_{h2} , respectively. For sufficient and unequal external activation signals, the T cell adopts the phenotype that is most strongly activated by the external signal. When the cell is externally stimulated by a T_{h1} signal, the naive state disappears (Fig. 5b). As a consequence, the former location of the naive state falls within the basin of attraction of the T_{h1} (20) state (Fig. 5e). The cell is therefore driven towards the T_{h1} phenotype in response to a T_{h1} signal. Analogously, a T_{h2} signal drives a cell towards the T_{h2} phenotype. In response to increasing T_{h1} stimulation, the basin of attraction of the T_{h2} state shrinks and ultimately the T_{h2} state disappears (Fig. 5c and f). If the cell is in one of the states that is eliminated, i.e. state 00 or 02, it will go to the only stable state left which is state 20 (Fig. 5f). Effectively, helper T cell skewing blocks the cell from adopting the opposite phenotype by temporarily eliminating the other stable states from the system for the duration of the external stimulus.

An n -dimensional model

The extended framework for helper T cell differentiation currently comprises many putative master regulators that control different T helper phenotypes. When regulation of other master regulators, such as ROR γ t and FoxP3, is compared with that of Tbet and GATA3, many parallels emerge. All four are activated by NFAT and different components of the

Jak-Stat pathway: ROR γ t is up-regulated by Stat3 (35) and FoxP3 by Stat3 and Stat5 (36, 37). Despite the partial overlap in specific inducing cytokine and Jak-Stat signals, the expression of all four master regulators remains mutually exclusive with respect to each other. Intracellular self-activation has not been demonstrated for ROR γ t and FoxP3, but both reinforce their own expression at least via the extracellular cytokine route (7, 38).

To accommodate these master regulators, the two-dimensional master regulator model can be expressed as an n -dimensional framework. The maximum number of steady states depends on the total number of master regulators (i.e. n) in the model: three steady states per master regulator (resting, saddle and active) allows for a maximum of 3^n possible steady states, of which 2^n states are stable. All genes are transcribed and translated by the same molecular machinery; hence, all genes are assigned the same productivity ρ , making the total number of parameters in such a model is $1 + 2n + n^2$.

To model each of these master regulators, they are each denoted by M_i for $i = 1$ to n :

$$\frac{dM_i}{d\tau} = \rho \left[\frac{\alpha_i M_i^2 + \varepsilon_i}{1 + \alpha_i M_i^2 + \varepsilon_i} \right] - M_i \left(1 + \sum_{j=1}^n \mathbf{B}_{ij} M_j \right) \quad (10)$$

where α_i is the self-activation parameter, ε_i is the external activation parameter and \mathbf{B}_{ij} is a matrix of complex formation parameters. This notation conveniently integrates parameter




Fig. 5. Phase plane analysis of the T_{h1}/T_{h2} (two-dimensional) master regulator model. The upper panels (a, b and c) show nullclines and steady states, with the corresponding separatrixes (panels d and e) or exemplary trajectories (panel f, grey lines) shown below. The black lines are the $\frac{dM_2}{dt} = 0$ nullclines and the grey those of M_1 (panels a, b and c). The naming convention for states was chosen such that the four stable states (solid squares) have only even numbers (0 and 2) in their state name, whereas the five non-stable states (open squares) also contain a 1 in their state name. The vector field is indicated by arrows in each qualitatively different area. Parameter values: $\rho = 5$, $\beta = 1$, $\mu_1 = \mu_2 = \alpha_1 = \alpha_2 = 1$ and $\varepsilon_2 = 0.01$. In panels a and d, we set $\varepsilon_1 = 0.01$ to show all possible steady states; in panels b and e, we set $\varepsilon_1 = 0.02$ (mild T_{h1} skewing) and $\beta = 1$ and in panels c and f, we set $\varepsilon_1 = 0.1$ (strong T_{h1} skewing) to show the effect of asymmetric external stimulation.

μ_i and β_{ij} into a single matrix; the homodimerization parameters μ_i are the diagonal of this matrix, and inhibition parameters β_{ij} form all off-diagonal elements. As yet, there is hardly any information about the exact topology of this matrix, but this is expected to change in the near future.

Accounting for FoxP3/GATA3 data

To illustrate the application of the model to data, we searched for parameter values that describe master regulator kinetics in human CD4+CD45RA+ T cells. These data, published by Mantel *et al.* (22), comprise FoxP3 and GATA3 mRNA expression levels measured in naive T_h after stimulation under either Treg-skewing conditions (plate bound aCD3-aCD28 + TGF β 5 ng ml $^{-1}$), T_h2 skewing (plate bound aCD3-aCD28 + IL-4, 25 ng ml $^{-1}$) or T_h0 (stimulation with plate bound aCD3-aCD28 only). FoxP3 and GATA3 expressions were measured at 0, 2, 6, 24 and 96 h after stimulation. For details concerning the data and the exact

experimental set-up, we refer to the original publication by Mantel *et al.* (22).

Taking the limited number of available data points and the relatively large number of model parameter into account, we searched manually for an illustrative fit and refrained from performing a full non-linear parameter fit. To mark the transition from a resting to an active cell, we let gene production ρ increase over time (see the legend of Fig. 6), as naive T cells have been shown to slowly increase their rates of transcription and translation in response to activation (39). Increasing the gene production parameter ρ facilitates the cell to approach the high-expression steady states (Fig. 3a).

The kinetics of master regulators FoxP3 and GATA3 in this system are described reasonably well by our model (Fig. 6). The model clearly accounts for the establishment of a dominant master regulator under certain polarizing conditions: FoxP3 for Treg conditions (Fig. 6a) and GATA3 for T_h2 conditions (Fig. 6b). Furthermore, for the same parameter




Fig. 6. The master regulator model agrees with experimental data of Mantel *et al.* (22). T_h were stimulated under three different conditions: Treg (a), T_h2 (b) and T_h0 (c). FoxP3 (M_1) and GATA3 measurements (M_2) are depicted as squares and circles, respectively, with error bars indicated. For visualization reasons, the data are plotted slightly offset with respect to time; all data were collected at 0, 2, 6, 24 or 96 h. In (d), increasing IL-4 concentration is modelled using the parameter ε_2 as a bifurcation parameter, and the data are plotted similarly as in (a–c). Parameter values: $\rho = 0.3 + \frac{1.75t}{t+30}$, where t is time (hours), $\mu_1 = \mu_2 = 0.01$, $\alpha_1 = 4.5$, $\alpha_2 = 4.0$ and inhibition $\beta = 2$. In panel a, we set $\varepsilon_1 = 1$ and $\varepsilon_2 = 0.25$; in panel b, $\varepsilon_1 = 0.4$ and $\varepsilon_2 = 2$; in panel c, we set $\varepsilon_1 = 0.5$ and $\varepsilon_2 = 0.5$. In panel d, we set ρ to 1.7 (i.e. we fix time at $t = 120$) in accordance with the experiment readout time of 5 days; $\varepsilon_1 = 0.6$, $\varepsilon_2 = [\text{IL-4 (pg ml}^{-1})]/2$.

values, the model describes the dynamics of these master regulators stimulated only with activation signals, i.e. in a non-polarizing T_h0 environment (Fig. 6c). In addition to describing the dynamics of FoxP3 and GATA3 in time course experiments, the model accounts for the response of naive T cells to conflicting differentiation signals. Naive T_h were activated and stimulated with TGF β and different concentrations of IL-4. Expressions of FoxP3 and GATA3 were quantified after 5 days of stimulation. To implement these conditions in the model, the external activator IL-4 (parameter ε_2) was varied and the model was observed at $t = 120$, i.e. after 5 days. The model should not be expected to fit this data perfectly, because it remains a caricature model missing (for instance) regulatory elements in the signal transduction pathway that connects extracellular IL-4 concentration to GATA3 expression.

Discussion

The mathematical model presented here describes the differentiation of helper T cells by the expression of master regulators, which are transcription factors that are both necessary and sufficient for the induction of a certain cellular phenotype (4). We have shown that a bistable master regulator can be switched from a state of low to high expression by extracellular cytokine stimulation. Co-expression of master regulators can be prevented by strong mutual inhibition through complex formation. Transient helper T cell skewing prevents the cell from differentiating towards the opposite phenotype by temporarily eliminating the opposite stable state from the system. Once polarized, the cell resists phenotype switching and hence displays phenotype memory. These properties are sufficient to explain the differentiation of a T cell into several exclusive phenotypes.

Previous models have described only one (18) or two (19–21, 40) master regulators. Our model differs from these models by using a more mechanistic rather than a phenomenological description of transcription factor binding. Firstly, using dimerization of transcription factors and binding to DNA, we show that the activation term can be approximated by Michaelis–Menten kinetics describing enzyme–multiple substrate interactions. Furthermore, mutual inhibition is described mechanistically by master regulator sequestration into a nuclear complex rather than a generic inhibition term. Despite the fact that our model is more mechanistic, it is also more simple than earlier models, and yet we obtain similar results in one and two dimensions. The ability of our extended model to describe any number of master regulators allows for more complex networks of genes to be analysed and permits incorporation of newly discovered phenotypes and their master regulators.

To illustrate the model's ability to describe real data, we show that the model can account for data published by Mantel *et al.* (22). Experiments that would allow us to fit this model to data are rare, because studies typically perform measurements at one or two time points or provide qualitative rather than quantitative experimental readouts. To our knowledge, this is the first attempt to describe helper T cell transcription factors over time. The model that we present disregards much of the known detail about induction and propagation of master regulator genes, but the agreement between model and data suggests that we have may succeeded in capturing the essence of the regulation of these

genes in this caricature model. Strictly speaking, a model is always incorrect, because it only approximates reality. However, a good model allows one to focus on the important processes that primarily determine master regulator dynamics. In other words, the model developed here is 'no more, but no less, than a way of thinking clearly' (34).

Further confirmation and testing of the model would need detailed data about master regulator kinetics in time. Using a data set that measures all master regulators simultaneously in response to different (and if possible, conflicting) stimuli, one could estimate master regulator-specific parameters. Although a good fit to experimental data by no means proves that a model is accurate, it is a minimal requirement for a correct model. As a test of the framework of the model, additional biochemical studies into the mechanism of inhibition between the master regulators are required. Indeed, it was shown that retrovirus-mediated up-regulation Tbet or GATA3 in IL-4 and IFN γ knockout mice is sufficient to inhibit FoxP3 (41), which is consistent with the inhibition by complex formation. Furthermore, an interaction between FoxP3 and ROR γ t was recently shown to be an inhibitory mechanism in T_h differentiation (42).

The application of this model is not limited to T cell biology or even immunology. The generic features described in the model can be found in many other developmental systems. Indeed, the master regulator has been adopted for describing developmental processes in general (43, 44), and this mathematical model could similarly be applied to these gene regulatory networks involved in cellular differentiation.

Funding

VIRGO consortium, an Innovative Cluster approved by The Netherlands Genomics Initiative; Dutch Government (BSIK 03012); The Netherlands Organisation for Scientific Research (NWO) VICI (016.048.603).

Abbreviation

TGF	transforming growth factor
NF κ B	nuclear factor κ B
NFAT	nuclear factor of activated T cells

References

- Mosmann, T. R., Cherwinski, H., Bond, M. W., Giedlin, M. A. and Coffman, R. L. 1986. Two types of murine helper T cell clone. I. Definition according to profiles of lymphokine activities and secreted proteins. *J. Immunol.* 136:2348.
- Shier, P., Hofstra, C. L., Ma, X. J. *et al.* 2000. Tbt-1, a new T-box transcription factor induced in activated Th1 and CD8 $^+$ T cells. *Immunogenetics* 51:771.
- Ferber, I. A., Lee, H. J., Zonin, F. *et al.* 999. GATA-3 significantly downregulates IFN-gamma production from developing Th1 cells in addition to inducing IL-4 and IL-5 levels. *Clin. Immunol.* 91:134.
- Rothenberg, E. V. 2007. Cell lineage regulators in B and T cell development. *Nat. Immunol.* 8:441.
- Kelso, A., Trout, A. B., Maraskovsky, E. *et al.* 1991. Heterogeneity in lymphokine profiles of CD4 $^+$ and CD8 $^+$ T cells and clones activated *in vivo* and *in vitro*. *Immunol. Rev.* 123:85.
- Ley, K., Smith, E. and Stark, M. A. 2006. IL-17A-producing neutrophil-regulatory Tn lymphocytes. *Immunol. Res.* 34:229.
- Weaver, C. T., Harrington, L. E., Mangan, P. R., Gavioli, M. and Murphy, K. M. 2006. Th17: an effector CD4 T cell lineage with regulatory T cell ties. *Immunity* 24:677.

- 8 Bettelli, E., Oukka, M. and Kuchroo, V. K. 2007. T(H)-17 cells in the circle of immunity and autoimmunity. *Nat. Immunol.* 8:345.
- 9 Veldhoen, M., Hocking, R. J., Atkins, C. J., Locksley, R. M. and Stockinger, B. 2006. TGFbeta in the context of an inflammatory cytokine milieu supports *de novo* differentiation of IL-17-producing T cells. *Immunity* 24:179.
- 10 Ivanov, I. I., McKenzie, B. S., Zhou, L. *et al.* 2006. The orphan nuclear receptor RORgamma directs the differentiation program of proinflammatory IL-17⁺ T helper cells. *Cell* 126:1121.
- 11 Yang, X. O., Pappu, B. P., Nurieva, R. *et al.* 2008. T helper 17 lineage differentiation is programmed by orphan nuclear receptors RORalpha and RORgamma. *Immunity* 28:29.
- 12 Acosta-Rodriguez, E. V., Napolitani, G., Lanzavecchia, A. and Sallusto, F. 2007. Interleukins 1beta and 6 but not transforming growth factor-beta are essential for the differentiation of interleukin 17-producing human T helper cells. *Nat. Immunol.* 8:942.
- 13 Chen, Z., Tato, C. M., Muul, L., Laurence, A. and O'Shea, J. J. 2007. Distinct regulation of interleukin-17 in human T helper lymphocytes. *Arthritis. Rheum.* 56:2936.
- 14 Li, M. O., Wan, Y. Y., Sanjabi, S., Robertson, A. K. and Flavell, R. A. 2006. Transforming growth factor-beta regulation of immune responses. *Annu. Rev. Immunol.* 24:99.
- 15 Fontenot, J. D., Rasmussen, J. P., Williams, L. M., Dooley, J. L., Farr, A. G. and Rudensky, A. Y. 2005. Regulatory T cell lineage specification by the forkhead transcription factor foxp3. *Immunity* 22:329.
- 16 Williams, L. M. and Rudensky, A. Y. 2007. Maintenance of the Foxp3-dependent developmental program in mature regulatory T cells requires continued expression of Foxp3. *Nat. Immunol.* 8:277.
- 17 Deenick, E. K. and Tangye, S. G. 2007. Autoimmunity: IL-21: a new player in Th17-cell differentiation. *Immunol. Cell Biol.* 85:503.
- 18 Hofer, T., Nathansen, H., Lohning, M., Radbruch, A. and Heinrich, R. 2002. GATA-3 transcriptional imprinting in Th2 lymphocytes: a mathematical model. *Proc. Natl Acad. Sci. USA* 99:9364.
- 19 Mariani, L., Lohning, M., Radbruch, A. and Hofer, T. 2004. Transcriptional control networks of cell differentiation: insights from helper T lymphocytes. *Prog. Biophys. Mol. Biol.* 86:45.
- 20 Yates, A., Callard, R. and Stark, J. 2004. Combining cytokine signalling with T-bet and GATA-3 regulation in Th1 and Th2 differentiation: a model for cellular decision-making. *J. Theor. Biol.* 231:181.
- 21 Callard, R. E. 2007. Decision-making by the immune response. *Immunol. Cell Biol.* 85:300.
- 22 Mantel, P. Y., Kuipers, H., Boyman, O. *et al.* 2007. GATA3-driven Th2 responses inhibit TGF-beta1-induced FOXP3 expression and the formation of regulatory T cells. *PLoS Biol.* 5:e329.
- 23 Scheffer, M., Carpenter, S., Foley, J. A., Folke, C. and Walker, B. 2001. Catastrophic shifts in ecosystems. *Nature* 413:591.
- 24 Murphy, K. M. and Reiner, S. L. 2002. The lineage decisions of helper T cells. *Nat. Rev. Immunol.* 2:933.
- 25 Ansel, K. M., Djuretic, I., Tanasa, B. and Rao, A. 2006. Regulation of Th2 differentiation and Il4 locus accessibility. *Annu. Rev. Immunol.* 24:607.
- 26 Macian, F. 2005. NFAT proteins: key regulators of T-cell development and function. *Nat. Rev. Immunol.* 5:472.
- 27 Mowen, K. A. and Glimcher, L. H. 2004. Signaling pathways in Th2 development. *Immunol. Rev.* 202:203.
- 28 Ouyang, W., Lohning, M., Gao, Z. *et al.* 2000. Stat6-independent GATA-3 autoactivation directs IL-4-independent Th2 development and commitment. *Immunity* 12:27.
- 29 Mullen, A. C., Hutchins, A. S., High, F. A. *et al.* 2002. Hlx is induced by and genetically interacts with T-bet to promote heritable T(H)1 gene induction. *Nat. Immunol.* 3:652.
- 30 Afkarian, M., Sedy, J. R., Yang, J. *et al.* 2002. T-bet is a STAT1-induced regulator of IL-12R expression in naive CD4⁺ T cells. *Nat. Immunol.* 3:549.
- 31 Zhu, J., Yamane, H., Cote-Sierra, J., Guo, L. and Paul, W. E. 2006. GATA-3 promotes Th2 responses through three different mechanisms: induction of Th2 cytokine production, selective growth of Th2 cells and inhibition of Th1 cell-specific factors. *Cell Res.* 16:3.
- 32 Hwang, E. S., Szabo, S. J., Schwartzberg, P. L. and Glimcher, L. H. 2005. T helper cell fate specified by kinase-mediated interaction of T-bet with GATA-3. *Science* 307:430.
- 33 Chen, G. Y., Osada, H., Santamaria-Babi, L. F. and Kannagi, R. 2006. Interaction of GATA-3/T-bet transcription factors regulates expression of sialyl Lewis X homing receptors on Th1/Th2 lymphocytes. *Proc. Natl Acad. Sci. USA* 103:16894.
- 34 May, R. M. 2004. Uses and abuses of mathematics in biology. *Science* 303:790.
- 35 Yang, X. O., Panopoulos, A. D., Nurieva, R. *et al.* 2007. STAT3 regulates cytokine-mediated generation of inflammatory helper T cells. *J. Biol. Chem.* 282:9358.
- 36 Zorn, E., Nelson, E. A., Mohseni, M. *et al.* 2006. IL-2 regulates FOXP3 expression in human CD4⁺CD25⁺ regulatory T cells through a STAT-dependent mechanism and induces the expansion of these cells *in vivo*. *Blood* 108:1571.
- 37 Chen, Z., Laurence, A. and O'Shea, J. J. 2007. Signal transduction pathways and transcriptional regulation in the control of Th17 differentiation. *Semin. Immunol.* 19:400.
- 38 Bettelli, E., Korn, T. and Kuchroo, V. K. 2007. Th17: the third member of the effector T cell trilogy. *Curr. Opin. Immunol.* 19:652.
- 39 Hamalainen, H. K., Tubman, J. C., Vikman, S. *et al.* 2001. Identification and validation of endogenous reference genes for expression profiling of T helper cell differentiation by quantitative real-time RT-PCR. *Anal. Biochem.* 299:63.
- 40 Yates, A., Bergmann, C., Van Hemmen, J. L., Stark, J. and Callard, R. 2000. Cytokine-modulated regulation of helper T cell populations. *J. Theor. Biol.* 206:539.
- 41 Wei, J., Duramad, O., Perng, O. A., Reiner, S. L., Liu, Y. J. and Qin, F. X. 2007. Antagonistic nature of T helper 1/2 developmental programs in opposing peripheral induction of Foxp3⁺ regulatory T cells. *Proc. Natl Acad. Sci. USA* 104:18169.
- 42 Zhou, L., Lopes, J. E., Chong, M. M. *et al.* 2008. TGF-beta-induced Foxp3 inhibits T(H)17 cell differentiation by antagonizing RORgamma function. *Nature* 453:236.
- 43 O'Malley, B. W. 2007. Coregulators: from whence came these "master genes". *Mol. Endocrinol.* 21:1009.
- 44 Garraway, L. A. and Sellers, W. R. 2006. Lineage dependency and lineage-survival oncogenes in human cancer. *Nat. Rev. Cancer* 6:593.