

Horizontal gene transfer can rescue prokaryotes from Muller's ratchet

N. Takeuchi¹, K. Kaneko¹, E.V. Koonin²

¹ University of Tokyo, Japan

² National Center for Biotechnology Information, USA

Society for General Microbiology Annual Conference
April 16, 2014, Liverpool

Evolutionary significance of genetic recombination

- has been extensively considered in relation to evolution of sexual reproduction in eukaryotes
- But, many prokaryotes also undergo recombination → known as horizontal gene transfer (HGT)
 - How frequent is HGT? → it can be more freq. than mutation
 - S. pneumoniae, N. meningitidis* (Feil et al '00):
 - Event ratio: HGT/mutation \approx 5-10
 - Base substitution ratio: HGT/mutation \approx 100-50
 - Other various prokaryotes (Vos & Didelot '09):
 - Base substitution ratio: HGT/mutation \approx 100-50
- Is exchange of genetic information important for life in general?


HGT is different from meiotic recombination

○ Mechanism of HGT

- Transformation ←
- Transduction
- Conjugation

○ Transformation


Cells absorb DNA in environment and undergo recombination




Exchange of genetic information happens indirectly via environment

Muller's ratchet

- Assumption: Genome has undergone sufficient evolutionary adaptation
→ Mutations always decrease fitness



- Frequency distribution of mutations per genome (Haigh '78)




- Mutations accumulate in finite population (Muller '64; Felsenstein '74)


Relationship between Muller's ratchet and recombination

Case of eukaryotes (meiotic recombination)

- Recombination shuffles mutations between chromosomes




- Recombination can stop Muller's ratchet
(Pamilo et al. '87; Charlesworth & Charlesworth '93)




How Muller's ratchet is related to recombination

Case of prokaryotes (HGT)

- Recombination shuffles mutations via environmental DNA (eDNA)




- Deleterious mutations increase mortality of cells, so eDNA on average contains more mutations than the population





Model

- Model = population + eDNA pool
- Population dynamics
 - Wright-Fisher process
 - mutation + eDNA uptake (HGT)
 - fitness $f_i = (1-s)^i$
- eDNA pool dynamics
 - Input from dead bacteria
Death rate $1-f_i/f_{\max}$
 - Decay




HGT can stop Muller's ratchet




Potential benefit of indirect genetic exchange

- Incorporate population subdivision into model




- Expectation

- Each subpopulation is more susceptible to Muller's ratchet
- But, ratchets operate independently
→ complementation between subpopulations



Population subdivision and “gene-sharing” help to maintain genetic information



Summary

- HGT facilitates evolutionary maintenance of genetic information even if it on average tends to increase deleterious mutations
- To maintain genomic information, it is more advantageous to partition individuals into multiple subpopulations and let them “**cross-reference**” each other’s genetic information

Acknowledgements

- Paulien Hogeweg
 - Alexander Lobkovsky
 - Nen Saito
 - Yuri Wolf
-
- Japan Society for the Promotion of Sciences
 - NIH Intramural Research Program